ELSEVIER

Neuroscience Letters 265 (1999) 223–226

Neuroscience Letters

Author Index to Volume 265

Abe, K., see Hayashi, T., 37
Abe, K., see Suzuki, H., 163
Alcaraz-Zubeldia, M., see Boll, M.-C., 155
Amenta, F., see Sabbatini, M., 5
Antoons, G., see D'Hooge, R., 83
Argiolas, A., see Melis, M.R., 171

Bagnoli, P., see Traina, G., 123
Baldoni, E., see Sabbatini, M., 5
Ballesteros, M.A., see Gallo, M., 57
Balse, E., Suhr, R., Haaf, A., Kelche, C., Jackisch, R., Cassel, J.-C., The potentiation of amphetamine-induced hyperlocomotion by fimbria-fornix lesions in rats is abolished by intrahippocampal grafts rich in serotonergic neurons, 79
Barber, L.A., see Frayer, S.M., 17
Barker, J.L., see Zhang, L., 25
Basile, A.S., Brichta, A.M., Harris, B.D., Morse, D., Coling, D., Skolnick, P., Dizocilpine attenuates streptomycin-induced vestibulotoxicity in rats, 71
Baumann, M., see Suopanki, J., 53
Beauvillain, J.-C., see Soulié, C., 147
Beekes, M., see McBride, P.A., 135
Benarroch, E.E., see Miller, S.M., 75
Blin, O., see Rihet, P., 143
Boll, M.-C., Sotelo, J., Otero, E., Alcaraz-Zubeldia, M., Rios, C., Reduced ferroxidase activity in the cerebrospinal fluid from patients with Parkinson's disease, 155
Boyadjian, A., Marin, L., Danion, F., Veering in human locomotion: the role of the effectors, 21
Braak, E., see Braak, H., 67
Braak, H., Sandmann-Keil, D., Gai, W., Braak, E., Extensive axonal Lewy neurites in Parkinson's disease: a novel pathological feature revealed by α-synuclein immunocytochemistry, 67
Brichta, A.M., see Basile, A.S., 71

Caberlotto, L., Jimenez, P., Overstreet, D.H., Hurd, Y.L., Mathé, A.A., Fuxe, K., Alterations in neuropeptide Y levels and Y1 binding sites in the Flinders Sensitive Line rats, a genetic animal model of depression, 191
Cadoni, A., see Sabbatini, M., 5
Caillet-Boudin, M.-L., see Soulié, C., 147
Cassel, J.-C., see Balse, E., 79
Cha, J.-H., see Lee, M.-Y., 107
Chang, Y.H., see Zhang, L., 25
Chartier-Harlin, M.-C., see Soulié, C., 147
Chen, Y.C., see Liu, I.-M., 183
Cheng, J.-T., see Liu, I.-M., 183
Chi, T.-C., see Liu, I.-M., 183
Cho, K.S., Xiao, Y.M., So, K.-F., Diao, Y.C., Chung, S.K., Synergistic effect of optic and peripheral nerve grafts on sprouting of axon-like processes of axotomized retinal ganglion cells in adult hamsters, 175
Chul Son, B., see Lee, M.-Y., 107
Chun, M.-H., see Lee, M.-Y., 107
Chung, S.K., see Cho, K.S., 175
Cole, J., see Scarchilli, K., 139
Coling, D., see Basile, A.S., 71

Danion, F., see Boyadjian, A., 21
de Beaurepaire, R., see Slisli, Y., 29
De Deyn, P.P., see D'Hooge, R., 83
Deecke, L., see Endl, W., 115
Delacourte, A., see Soulié, C., 147
D'Emilia, D.M., Lipton, S.A., Ratio of S-nitrosohomocyst(e)ine to homocyst(e)ine or other thiols determines neurotoxicity in rat cerebrocortical cultures, 103
Depaulis, A., see Deransart, C., 131
Deransart, C., Riban, V., Lê, B.-T., Hechler, V., Marescaux, C., Depaulis, A., Evidence for the involvement of the pallidum in the modulation of seizures in a genetic model of absence epilepsy in the rat, 131
D'Hooge, R., De Deyn, P.P., Van de Vijver, G., Antoons, G., Raes, A., Van Bogaert, P.P., Uraemic guanidino compounds inhibit γ-aminobutyric acid-evoked whole cell currents in mouse spinal cord neurones, 83
Diao, Y.C., see Cho, K.S., 175
Diedrich, S.L., see Romeo, R.D., 167
Dupont-Wallois, L., see Soulié, C., 147

Edström, A., see Hornfelt, M., 87
Ekström, P.A.R., see Hornfelt, M., 87
Endl, W., Walla, P., Lindinger, G., Deecke, L., Lang, W., Event-related potential correlates of false recognitions of faces, 115

Feldman, A.N., see Zhang, L., 25
Frayer, S.M., Barber, L.A., Vasko, M.R., Activation of protein kinase C enhances peptide release from rat spinal cord slices, 17
Fujii, T., see Koyama, T., 33
Fuxe, K., see Caberlotto, L., 191

Gai, W., see Braak, H., 67
Gallo, M., Marquez, S.L., Ballesteros, M.A., Maldonado, A., Functional blockade of the parabrachial area by tetrodotoxin disrupts the acquisition of conditioned taste aversion induced by motion-sickness in rats, 57
García-Ayllón, M., Gómez, J.L., Vidal, C.J., Amphiphilic properties of acetylcholinesterase monomers in mouse plasma, 211
Gauthier, G.M., see Scarchilli, K., 139
Gilbert, J.R., see Veinbergs, I., 218
Gómez, J.L., see García-Ayllón, M., 211
Goyal, P., see Zaheer, A., 203

PII: S0304-3940(99)00304-3

Haaf, A., see Balse, E., 79
Hájek, I., see Jelínková-Vondrašová, D., 111
Haltia, M., see Suopanki, J., 53
Hari, R., see Numminen, J., 119
Harris, B.D., see Basile, A.S., 71
Hasbroucq, T., see Rihet, P., 143
Hayakawa, K., Nakashima, S., Ito, Y., Mizuta, K., Miyata, H., Nozawa, Y., Increased expression of phospholipase D1 mRNA during cAMP- or NGF- induced differentiation in PC12 cells, 127
Hayashi, T., see Suzuki, H., 163
Hayashi, T., Warita, H., Abe, K., Itoyama, Y., Expression of cyclin-dependent kinase 5 and its activator p35 in rat brain after middle cerebral artery occlusion, 37
Hechler, V., see Deransart, C., 131
Hingtgen, J.N., see Hwang, B.H., 151
Hodges, P.W., Richardson, C.A., Transversus abdominis and the superficial abdominal muscles are controlled independently in a postural task, 91
Honda, T., see Murayama, O., 61
Hoover, B.R., Marshall, J.F., Population characteristics of preproenkephalin mRNA-containing neurons in the globus pallidus of the rat, 199
Hornfelt, M., Edström, A., Ekström, P.A.R., Upregulation of cytosolic phospholipase A_2 correlates with apoptosis in mouse superior cervical and dorsal root ganglia neurons, 87
Hu, Q., see Zhang, L., 25
Hurd, Y.L., see Caberlotto, L., 191
Hwang, B.H., Kunkler, P.E., Tarricone, B.J., Hingtgen, J.N., Nurnberger Jr., J.I., Stress-induced changes of norepinephrine uptake sites in the locus coeruleus of C57BL/6J and DBA/2J mice: a quantitative autoradiographic study using [^{3}H]-tomoxetine, 151

Illes, P., see Krügel, U., 49
Illnerová, H., see Jelínková-Vondrašová, D., 111
Inamura, S., see Nakagawa, T., 64
Isa, T., see Sasaki, S., 13
Iseki, E., Marui, W., Kosaka, K., Uéda, K., Frequent coexistence of Lewy bodies and neurofibrillary tangles in the same neurons of patients with diffuse Lewy body disease, 9
Ito, Y., see Hayakawa, K., 127
Itoyama, Y., see Hayashi, T., 37
Itoyama, Y., see Suzuki, H., 163
Iwatsubo, T., see Murayama, O., 61

Jackisch, R., see Balse, E., 79
Jaramillo, M., see Paavilainen, P., 179
Jelínková-Vondrašová, D., Hájek, I., Illnerová, H., Adjustment of the human circadian system to changes of the sleep schedule under dim light at home, 111
Jimenez, P., see Caberlotto, L., 191
Joëls, M., see Karten, Y.J.G., 41
Ju, W.-K., see Lee, M.-Y., 107

Kaneko, M., see Nakagawa, T., 64
Karten, Y.J.G., Slagter, E., Joëls, M., Effect of long-term elevated corticosteroid levels on field responses to synaptic stimulation, in the rat CA1 hippocampal area, 41
Kawashima, K., see Koyama, T., 33
Kelche, C., see Balse, E., 79
Ki Kang, J., see Lee, M.-Y., 107
Kim, W.-K., S-nitrosation ameliorates homocysteine-induced neurotoxicity and calcium responses in primary culture of rat cortical neurons, 99
Kimura, K., see Suzuki, H., 163
Kitagawa, H., see Suzuki, H., 163
Kittner, H., see Krügel, U., 49
Kojo, I., Rovamo, J., Illusory motion capture and release in rotating 3-D figure in humans, 45
Kosaka, K., see Iseki, E., 9
Koyama, T., Nakajima, Y., Fujii, T., Kawashima, K., Enhancement of cortical and hippocampal cholinergic neurotransmission through 5-HT1A receptor-mediated pathways by BAY × 3702 in freely moving rats, 33
Krügel, U., Kittner, H., Illes, P., Adenosine 5′-triphosphate-induced dopamine release in the rat nucleus accumbens in vivo, 49
Kunkler, P.E., see Hwang, B.H., 151

Lahjouji, F., see Zhang, L., 25
Lang, W., see Endl, W., 115
Lê, B.-T., see Deransart, C., 131
Lee, M.-Y., Ju, W.-K., Cha, J.-H., Chul Son, B., Chun, M.-H., Ki Kang, J., Park, C.K., Expression of vascular endothelial growth factor mRNA following transient forebrain ischemia in rats, 107
Lee, Y.-S., Yoon, B.-W., Roh, J.-K., Neuroprotective effects of lamotrigine enhanced by flunarizine in gerbil global ischemia, 215
Li, B.-S., see Zhang, L., 25
Li, W., see Zhang, L., 25
Lim, R., see Zaheer, A., 203
Lindholm, D., see Skoglösa, Y., 207
Lindinger, G., see Endl, W., 115
Lipton, S.A., see D'Emilia, D.M., 103
Liu, I.-M., Chi, T.-C., Chen, Y.C., Lu, F.-H., Cheng, J.-T., Activation of opioid μ-receptor by loperamide to lower plasma glucose in streptozotocin-induced diabetic rats, 183
Lu, F.-H., see Liu, I.-M., 183

Ma, W., see Zhang, L., 25
Maldonado, A., see Gallo, M., 57
Mallory, M., see Veinbergs, I., 218
Mante, M., see Veinbergs, I., 218
Marescaux, C., see Deransart, C., 131
Maric, D., see Zhang, L., 25
Marin, L., see Boyadjian, A., 21
Marquez, S.L., see Gallo, M., 57
Marshall, J.F., see Hoover, B.R., 199
Marui, W., see Iseki, E., 9
Masliah, E., see Veinbergs, I., 218
Mathé, A.A., see Caberlotto, L., 191
Mazarati, A.M., Wasterlain, C.G., *N*-methyl-D-asparate receptor antagonists abolish the maintenance phase of self-sustaining status epilepticus in rat, 187
McBride, P.A., Beekes, M., Pathological PrP is abundant in sympathetic and sensory ganglia of hamsters fed with scrapie, 135
Melis, M.R., Spano, M.S., Succu, S., Argiolas, A., The oxytocin antagonist d$(CH_2)_5$Tyr(Me)^2-Orn^8-vasotocin reduces non-contact penile erections in male rats, 171
Mestdagh, N., Wülfert, E., Bicuculline increases Ca^{2+} transients in rat cerebellar granule cells through non-$GABA_A$ receptor associated mechanisms, 95
Miller, S.M., Schmalz, P.F., Benarroch, E.E., Szurszewski, J.H., Leptin receptor immunoreactivity in sympathetic prevertebral ganglion neurons of mouse and rat, 75
Mitchell, V., see Soulié, C., 147
Miyata, H., see Hayakawa, K., 127
Mizugaki, M., see Suzuki, H., 163

Mizuta, K., see Hayakawa, K., 127
Morse, D., see Basile, A.S., 71
Murayama, M., see Murayama, O., 61
Murayama, O., Tomita, T., Nihonmatsu, N., Murayama, M., Sun, X., Honda, T., Iwatsubo, T., Takashima, A., Enhancement of amyloid β 42 secretion by 28 different presenilin 1 mutations of familial Alzheimer's disease, 61

Näätänen, R., see Paavilainen, P., 179
Nagano, S., Ogawa, Y., Yanagihara, T., Sakoda, S., Benefit of a combined treatment with trientine and ascorbate in familial amyotrophic lateral sclerosis model mice, 159
Naito, K., see Sasaki, S., 13
Nakagawa, T., Kaneko, M., Inamura, S., Satoh, M., Intracerebroventricular administration of nocistatin reduces inflammatory hyperalgesia in rats, 64
Nakajima, Y., see Koyama, T., 33
Nakashima, S., see Hayakawa, K., 127
Nihonmatsu, N., see Murayama, O., 61
Nozawa, Y., see Hayakawa, K., 127
Numminen, J., Salmelin, R., Hari, R., Subject's own speech reduces reactivity of the human auditory cortex, 119
Nurnberger Jr., J.I., see Hwang, B.H., 151

Ogawa, Y., see Nagano, S., 159
Otero, E., see Boll, M.-C., 155
Overstreet, D.H., see Caberlotto, L., 191

Paavilainen, P., Jaramillo, M., Näätänen, R., Winkler, I., Neuronal populations in the human brain extracting invariant relationships from acoustic variance, 179
Park, C.K., see Lee, M.-Y., 107
Patrone, C., see Skoglösa, Y., 207
Possamaï, C.-A., see Rihet, P., 143

Qiao, J.-T., see Wang, H., 195

Raes, A., see D'Hooge, R., 83
Riban, V., see Deransart, C., 131
Richardson, C.A., see Hodges, P.W., 91
Rihet, P., Hasbroucq, T., Blin, O., Possamaï, C.-A., Serotonin and human information processing: an electromyographic study of the effects of fluvoxamine on choice reaction time, 143
Rios, C., see Boll, M.-C., 155
Rockenstein, E., see Veinbergs, I., 218
Roh, J.-K., see Lee, Y.-S., 215
Romeo, R.D., Diedrich, S.L., Sisk, C.L., Estrogen receptor immunoreactivity in prepubertal and adult male Syrian hamsters, 167
Rovamo, J., see Kojo, I., 45
Rubinow, D.R., see Zhang, L., 25

Sabbatini, M., Baldoni, E., Cadoni, A., Vitaioli, L., Zicca, A., Amenta, F., Forebrain white matter in spontaneously hypertensive rats: a quantitative image analysis study, 5
Sakoda, S., see Nagano, S., 159
Salmelin, R., see Numminen, J., 119
Sandmann-Keil, D., see Braak, H., 67
Sasaki, S., Isa, T., Naito, K., Effects of lesion of pontomedullary reticular formation on visually triggered vertical and oblique head orienting movements in alert cats, 13
Satoh, M., see Nakagawa, T., 64
Scarchilli, K., Vercher, J.-L., Gauthier, G.M., Cole, J., Does the oculo-manual co-ordination control system use an internal model of the arm dynamics?, 139
Schmalz, P.F., see Miller, S.M., 75
Schubotz, R., Instruction differentiates the processing of temporal and spatial sequential patterns: evidence from slow wave activity in humans, 1
Sisk, C.L., see Romeo, R.D., 167
Skoglösa, Y., Patrone, C., Lindholm, D., Pituitary adenylate cyclase activating polypepetide is expressed by developing rat Purkinje cells and decreases the number of cerebellar γ-amino butyric acid positive neurons in culture, 207
Skolnick, P., see Basile, A.S., 71
Slagter, E., see Karten, Y.J.G., 41
Slisli, Y., de Beaurepaire, R., Interleukin-1β and calcitonin, but not corticotropin-releasing factor, alter sleep cycles when injected into the rat hypothalamic lateral paraventricular area, 29
So, K.-F., see Cho, K.S., 175
Sotelo, J., see Boll, M.-C., 155
Soulié, C., Mitchell, V., Dupont-Wallois, L., Chartier-Harlin, M.-C., Beauvillain, J.-C., Delacourte, A., Caillet-Boudin, M.-L., Synthesis of apolipoprotein E (ApoE) mRNA by human neuronal-type SK N SH-SY 5Y cells and its regulation by nerve growth factor and ApoE, 147
Spano, M.S., see Melis, M.R., 171
Succu, S., see Melis, M.R., 171
Suhr, R., see Balse, E., 79
Sun, X., see Murayama, O., 61
Suopanki, J., Tyynelä, J., Baumann, M., Haltia, M., Palmitoyl-protein thioesterase, an enzyme implicated in neurodegeneration, is localized in neurons and is developmentally regulated in rat brain, 53
Suzuki, H., Hayashi, T., Tojo, S.J., Kitagawa, H., Kimura, K., Mizugaki, M., Itoyama, Y., Abe, K., Anti-P-selectin antibody attenuates rat brain ischemic injury, 163
Szurszewski, J.H., see Miller, S.M., 75

Takashima, A., see Murayama, O., 61
Tarricone, B.J., see Hwang, B.H., 151
Tojo, S.J., see Suzuki, H., 163
Tomita, T., see Murayama, O., 61
Traina, G., Bagnoli, P., Mechanisms mediating somatostatin-induced reduction of cytosolic free calcium in PC12 cells, 123
Tyynelä, J., see Suopanki, J., 53

Uéda, K., see Iseki, E., 9

Van Bogaert, P.P., see D'Hooge, R., 83
Van de Vijver, G., see D'Hooge, R., 83
Vasko, M.R., see Frayer, S.M., 17
Veinbergs, I., Mallory, M., Mante, M., Rockenstein, E., Gilbert, J.R., Masliah, E., Differential neurotrophic effects of apolipoprotein E in aged transgenic mice, 218
Vercher, J.-L., see Scarchilli, K., 139
Vidal, C.J., see García-Ayllón, M., 211
Vitaioli, L., see Sabbatini, M., 5

Walla, P., see Endl, W., 115
Wang, H., Zhang, R.-x., Wang, R., Qiao, J.-T., Decreased expression of *N*-methyl-D-aspartate (NMDA) receptors in rat dorsal root ganglion following complete Freund's adjuvant-induced inflammation: an immunocytochemical study for NMDA NR1 subunit, 195
Wang, R., see Wang, H., 195

Warita, H., see Hayashi, T., 37
Wasterlain, C.G., see Mazarati, A.M., 187
Weiss, J.L., see Zaheer, A., 203
Winkler, I., see Paavilainen, P., 179
Wülfert, E., see Mestdagh, N., 95

Xiao, Y.M., see Cho, K.S., 175

Yanagihara, T., see Nagano, S., 159
Yoon, B.-W., see Lee, Y.-S., 215

Zaheer, A., Weiss, J.L., Goyal, P., Lim, R., Enhanced expression of neurotrophic factors by C6 rat glioma cells after transfection with glia maturation factor, 203
Zhang, L., Chang, Y.H., Feldman, A.N., Ma, W., Lahjouji, F., Barker, J.L., Hu, Q., Maric, D., Li, B.-S., Li, W., Rubinow, D.R., The expression of GABAA receptor α2subunit is upregulated by testosterone in rat cerebral cortex, 25
Zhang, R.-x., see Wang, H., 195
Zicca, A., see Sabbatini, M., 5